

F8

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C07D 307/00</b>		<b>A2</b>	(11) International Publication Number: <b>WO 00/69841</b>
			(43) International Publication Date: 23 November 2000 (23.11.00)
(21) International Application Number: <b>PCT/EP00/04015</b>		[GB/GB]; 2 Paternoster Court, Cassington Road Yarton, Oxford OX5 1QB (GB). BRIGGS, Barbara [GB/GB]; 24 Latchmere Road, Kingston-upon-Thames, Surrey KT2 5TW (GB). CONCEPTION, Arnel [PH/JP]; 3-6-27-207, Shibatsuji-cho, Nara City, Nara 630-8114 (JP). BULLOCK, William [US/US]; 60 Hermann Lane, Easton, CT 06612 (US).  (74) Common Representative: <b>BAYER AKTIENGESELLSCHAFT</b> ; D-51368 Leverkusen (DE).  (81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: <b>4 May 2000 (04.05.00)</b>			
(30) Priority Data: <b>9911453.0 17 May 1999 (17.05.99) GB</b>			
(71) Applicant (for all designated States except US): <b>BAYER AKTIENGESELLSCHAFT [DE/DE]; D-51368 Leverkusen (DE).</b>			
(72) Inventors; and (75) Inventors/Applicants (for US only): <b>BRÄUNLICH, Gabriele [DE/DE]; Claudiusweg 9, D-42115 Wuppertal (DE). ES-SAYED, Mazen [DE/DE]; Ricarda-Huch-Str. 36, D-40764 Langenfeld (DE). FISCHER, Rüdiger [DE/DE]; Zu den Fussfällen 23, D-50529 Pulheim (DE). FUGMANN, Burkhard [DE/DE]; Dr.-Kessel-Strasse 3, D-40878 Ratingen (DE). HENNING, Rolf [DE/DE]; Am Jagdhaus 115, D-42113 Wuppertal (DE). SCHNEIDER, Stephan [DE/DE]; Am Ringofen 46, D-42327 Wuppertal (DE). SPERZEL, Michael [DE/DE]; Normannenstrasse 31, D-42275 Wuppertal (DE). SCHLEMMER, Karl-Heinz [DE/DE]; Wildsteig 22a, D-42113 Wuppertal (DE). STURTON, Graham [GB/GB]; 184 Windsor Road, Bray Maidenhead SL6 2DW (GB). FITZGERALD, Mary</b>			
Published Without international search report and to be republished upon receipt of that report.			

(54) Title: BENZOFURANYLAMINOALCOHOLS



(57) Abstract

Benzofuranyl aminoalcohols of general formula (I), process for their preparation and their use in medicaments, especially for the treatment of inflammatory processes.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Benzofuranylaminoalcohols

The invention relates to Benzofuranylaminoalcohols, processes for their preparation and their use in medicaments.


5

It is known that the NADPH oxidase of phagocytes is the physiological source to the superoxide radical anion and reactive oxygen species derived therefrom which are important in the defence against pathogens. Moreover, both inflammatory (e.g.  $\text{TNF}\alpha$ , IL-1 or IL-6) and anti-inflammatory cytokines (e.g. IL-10) play a pivotal role in host defence mechanism. Uncontrolled production of inflammatory mediators can lead to acute or chronic inflammation, auto immune diseases, tissue damage, multi-organ failure and to death. It is additionally known that elevation of phagocyte cyclic AMP leads to inhibition of oxygen radical production and that this cell function is more sensitive than others such as aggregation or enzyme release.

15

Benzofuran derivatives having phosphodiesterase IV (PDE IV)-inhibiting action are described in EP 731 099. The reference describes only single-hydroxyl substituted derivatives, however. Benzofuranylaminoalcohols without 3-ureido moiety are disclosed for the treatment of diseases in the circulatory system in US 4,056,626. In order to provide alternative compounds with similar or improved PDE IV-inhibitory activity, the present invention relates to Benzofuranylaminoalcohols of the general formula (I)

20



25

in which

A represents hydrogen, straight-chain or branched acyl or alkoxycarbonyl, each having 1 to 6 carbon atoms, halogen, carboxyl, cyano, nitro, hydroxyl,

trifluoromethyl or trifluoromethoxy, or straight-chain or branched alkyl having 1 to 6 carbon atoms, which is optionally substituted by carboxyl, alkoxy or alkoxycarbonyl each having 1 to 4 carbon atoms, phenoxy or benzoyl,

5         $R^1$         represents hydrogen, straight-chain or branched alkyl having 1 to 4 carbon atoms, an amino protecting group or a group of the formula  $-CO-R^7$

in which

10         $R^7$         denotes straight chain or branched alkoxy having 1 to 4 carbon atoms,

$R^2$  and  $R^3$  are identical or different and represent hydrogen, cycloalkyl having 3, 4, 5 or 6 carbon atoms, straight chain or branched alkyl, alkoxycarbonyl or alkenyl each having 1 to 8 carbon atoms,

15

or

$R^2$  and  $R^3$  together with the nitrogen atom form a 5-, 6- or 7-membered saturated heterocycle optionally having a further oxygen atom,

20

$R^4$         represents aryl having 6 to 10 carbon atoms or  
represents a 5-, 6- or 7-membered, aromatic, saturated or unsaturated heterocycle, which can contain 1 to 3 oxygen, sulphur and/or nitrogen atoms as heteroatoms or a residue of a formula  $-NR^8$ ,

25

wherein

$R^8$         denotes hydrogen or straight-chain or branched alkyl or alkoxycarbonyl each having 1 to 6 carbon atoms,

30

and to which further a benzene ring can be fused and wherein aryl and/or the heterocycle are optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising hydroxyl, halogen, nitro, 1H-tetrazolyl, pyridyl, trifluoromethyl, trifluoromethoxy, difluoromethyl, difluoromethoxy, cyano, carboxy, straight-chain or branched alkoxy, alkoxycarbonyl or acyl each having 1 to 6 carbon atoms or by straight-chain or branched alkyl having 1 to 5 carbon atoms, which is optionally substituted by carboxyl or straight-chain or branched alkoxycarbonyl having 1 to 4 carbon atoms or by a group of formula  $-NR^9R^{10}$ ,  $-SR^{11}$ ,  $-(NH)_a-SO_2R^{12}$  or  $-O-SO_2R^{13}$ ,

in which

$R^9$  and  $R^{10}$  are identical or different and denote hydrogen or a straight-chain or branched alkyl having 1 to 4 carbon atoms,

or

$R^9$  denotes hydrogen

and

$R^{10}$  denotes straight-chain or branched acyl having 1 to 6 carbon atoms,

$R^{11}$  denotes hydrogen or straight-chain or branched alkyl having 1 to 4 carbon atoms,

$a$  denotes a number 0 or 1,

$R^{12}$  and  $R^{13}$  are identical or different and represent straight-chain or branched alkyl having 1 to 6 carbon atoms, benzyl or phenyl, which are optionally

substituted by trifluoromethyl, halogen or straight-chain or branched alkyl having 1 to 4 carbon atoms,

L represents an oxygen or sulfur atom

5

R<sup>5</sup> and R<sup>6</sup> represent hydrogen, straight-chain or branched alkyl having 1 to 6 carbon atoms, which is optionally substituted by aryl having 6 to 10 carbon atoms or by a 5- to 7-membered aromatic, saturated or unsaturated heterocycle having 1 to 3 heteroatoms from the series comprising N, S, O and/or a residue of a formula -NR<sup>14</sup>

10

wherein

R<sup>14</sup> denotes hydrogen or straight-chain or branched alkyl or alkoxycarbonyl each having 1 to 6 carbon atoms

15

and to which a phenyl ring can be fused and which are optionally monosubstituted to disubstituted by identical or different substituents from the series comprising halogen, cyano or by a straight-chain or branched alkyl or alkoxy-carbonyl each having 1 to 6 carbon atoms,

20

or represent a 6-membered saturated N-heterocycle, which is optionally substituted by alkoxycarbonyl having 1 to 6 carbon atoms,

25

or

R<sup>5</sup> and R<sup>6</sup> together with the nitrogen atom form a 5- to 6-membered aromatic, saturated or unsaturated heterocycle having 1 to 3 heteroatoms from the series comprising N, S, O and/or a residue of a formula -NR<sup>14</sup>,

30

and to which a phenyl ring can be fused and which is optionally monosubstituted to disubstituted by identical or different substituents from the series comprising halogen, cyano or by a straight-chain or branched alkyl or alkoxy-carbonyl each having 1 to 6 carbon atoms,

and salts thereof.

The benzofuranylaminoalcohols according to the invention can also be present in the form of their salts. In general, salts with organic or inorganic bases or acids may be mentioned here.

Physiologically acceptable salts are preferred in the context of the present invention. Physiologically acceptable salts of the benzofuranylaminoalcohols can be metal or ammonium salts of the substances according to the invention, which contain a free carboxylic group. Those which are particularly preferred are, for example, sodium, potassium, magnesium or calcium salts, and also ammonium salts which are derived from ammonia, or organic amines, such as, for example, ethylamine, di- or triethylamine, di- or triethanolamine, dicyclohexylamine, dimethylaminoethanol, arginine, lysine or ethylenediamine.

Physiologically acceptable salts can also be salts of the compounds according to the invention with inorganic or organic acids. Preferred salts here are those with inorganic acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid or sulphuric acid, or salts with organic carboxylic or sulphonic acids such as, for example, acetic acid, maleic acid, fumaric acid, malic acid, citric acid, tartaric acid, ethanesulphonic acid, benzenesulphonic acid, toluenesulphonic acid or naphthalenedisulphonic acid.

The compounds according to the invention can exist in stereoisomeric forms which either behave as image and mirror image (enantiomers), or which do not behave as

image and mirror image (diastereomers). The invention relates both to the antipodes and to the racemate forms, as well as to individual diastereomers and to the diastereomer mixtures. The racemate forms, like the diastereomers, can be separated into the stereoisomerically uniform constituents in a known manner.

5

Heterocycle in general represents a 5- to 7-membered aromatic, saturated or unsaturated, preferably 5- to 6- membered, aromatic or saturated ring, which can contain up to 3 oxygen, sulphur, nitrogen atoms or a residue of a formula  $-NR^{14}$  as heteroatoms, wherein  $R^{14}$  has the abovementioned meaning, and to which further aromatic rings can be fused.

10

The following are mentioned as preferred: thienyl, furyl, pyrrolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, quinolyl, isoquinolyl, quinazolyl, quinoxazolyl, thiazolyl, dihydrothiazolyl, benzothiazolyl, isothiazolyl, benzisothiazolyl, oxazolyl, benzoxa-

15

zolyl, isoxazolyl, imidazolyl, benzimidazolyl, indolyl, morpholinyl, pyrrolidinyl, piperidyl, piperazinyl, oxazolinyl or triazolyl.

Preferred compounds of the general formula (I) are those

20

in which

A represents hydrogen, halogen, carboxyl, cyano, nitro, hydroxyl, trifluoromethyl, trifluoromethoxy or alkoxy having up to 4 carbon atoms

25

$R^1$  represents hydrogen, straight-chain or branched alkyl having up to 4 carbon atoms or a group of the formula  $-CO-R^7$

in which

30

$R^7$  denotes straight chain or branched alkoxy having up to 4 carbon atoms,



R<sup>2</sup> and R<sup>3</sup> are identical or different and represent hydrogen, cyclobutyl, cyclopentyl, cyclohexyl or straight-chain or branched alkyl, alkoxycarbonyl or alkenyl each having up to 4 carbon atoms, or

5 or

R<sup>2</sup> and R<sup>3</sup> together with the nitrogen atom form a pyrrolidinyl-, piperidinyl- or morpholinyl-ring,

10 and

R<sup>4</sup> represents phenyl, pyridyl or thienyl, wherein all rings are optionally mono-substituted to trisubstituted by identical or different substituents from the series comprising hydroxyl, fluorine, chlorine, bromine, nitro, carboxy, straight-chain or branched alkoxy, alkoxycarbonyl or acyl each having up to 3 carbon atoms, or by straight-chain or branched alkyl having up to 3 carbon atoms, which is optionally substituted by carboxyl or straight-chain or branched alkoxycarbonyl having up to 3 carbon atoms

20 L represents an oxygen or sulfur atom,

R<sup>5</sup> and R<sup>6</sup> represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms, which is optionally substituted by phenyl, pyridyl, imidazolyl, pyrrol, morpholinyl, piperidinyl, piperazinyl, pyrrolidinyl, wherein the heterocycles optionally contain a residue of a formula -NR<sup>14</sup>,

25

wherein

R<sup>14</sup> denotes hydrogen or straight-chain or branched alkyl or alkoxycarbonyl each having up to 4 carbon atoms,

30

and wherein the ring systems are optionally monosubstituted by straight-chain or branched alkyl, alkoxy or alkoxycarbonyl each having up to 3 carbon atoms,

5 or represent a 6-membered saturated N-heterocycle, which is optionally substituted by alkoxycarbonyl having 1 to 6 carbon atoms,

or

10 R<sup>5</sup> and R<sup>6</sup> together with the nitrogen atom form a pyrazolyl-, triazolyl-, tetrazolyl-, imidazolyl-, pyrrol-, morpholinyl-, piperidinyl-, pyrrolidinyl-, piperazinylring, wherein the heterocycles optionally contain a residue of a formula -NR<sup>14</sup>,

15 and wherein the ringsystem is optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising halogen, cyano or by a straight-chain or branched alkyl having up to 6 carbon atoms,

and salts thereof.

20 Particularly preferred compounds of the general formula (I) are those in which

A represents hydrogen,

25 R<sup>1</sup> represents hydrogen or straight-chain or branched alkyl having up to 3 carbon atoms or a group of the formula -CO-R<sup>7</sup>,

in which

30 R<sup>7</sup> denotes straight chain or branched alkoxy having up to 3 carbon atoms,

R<sup>2</sup> and R<sup>3</sup> represent hydrogen,

5 R<sup>4</sup> represents phenyl or pyridyl, which are optionally up to difold substituted by identical or different substituents from the series fluorine, chlorine, methyl or methoxy,

L represents an oxygen atom,

10 R<sup>5</sup> and R<sup>6</sup> represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms, which is optionally substituted by phenyl, pyridyl, imidazolyl, pyrrol, piperidinyl, wherein the heterocycles optionally contain a residue of a formula -NR<sup>14</sup>,

15 wherein

R<sup>14</sup> denotes hydrogen or straight-chain or branched alkyl or alkoxycarbonyl each having up to 3 carbon atoms,

20 and wherein the ring systems are optionally monosubstituted by straight-chain or branched alkyl, alkoxy or alkoxycarbonyl each having up to 3 carbon atoms,

or represent a 6-membered saturated N-heterocycle, which is optionally substituted by alkoxycarbonyl having 1 to 6 carbon atoms,

25 or

R<sup>5</sup> and R<sup>6</sup> together with the nitrogen atom form a imidazolyl-, pyrrol-, morpholinyl-, piperidinyl-, pyrrolidinyl-, piperazinylring, wherein the heterocycles optionally contain a residue of a formula -NR<sup>14</sup>,

30

wherein

$R^{14}$  have the abovementioned meaning of  $R^{14}$  and is identical or different to the latter,



5 and wherein the ringsystem are optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising by a straight-chain or branched alkyl or alkoxycarbonyl each having up to 3 carbon atoms,

and salts thereof.




10

Very particular preferred compounds of the general formula (I) are those shown in table A:





Table A

Structure



## Structure




## Structure



A process for the preparation of the compounds of the general formula (I) has additionally been found, characterized in that

compounds of the general formula (II)

5



in which

A, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and L have the abovementioned meaning,

10

are reacted with amines of the general formula (III)



15


in which

R<sup>5</sup> and R<sup>6</sup> have the abovementioned meaning,

in the presence of an inert solvent.

20

The process according to the invention can be illustrated by way of example by the following equations:



5 Suitable solvents are generally customary organic solvents which do not change under the reaction conditions. These include ethers such as diethyl ether, dioxane or tetrahydrofuran, acetone, dimethylsulfoxide, dimethylformamide or alcohols such as methanol, ethanol, propanol or halogenohydrocarbons such as dichloromethane, trichloromethane or tetrachloromethane and mixtures of two or three of the abovementioned solvents. A mixture of methanol and tetrahydrofuran is preferred.


10 The process is in general carried out in a temperature range from  $-30^\circ\text{C}$  to  $+100^\circ\text{C}$ , preferably from  $0^\circ\text{C}$  to  $50^\circ\text{C}$ .

The process is generally carried out at normal pressure. However, it is also possible to carry out it at elevated or reduced pressure (for example in a range from 0.5 to 5 bar).

15

The compounds of the general formula (II) are new and can be prepared by reacting compounds of the general formula (IV)






in which,

L, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> have the abovementioned meaning,

5

with glycidol of the formula (V)



10 in the presence of triphenylphosphin and diethylazodicarboxylate in an inert solvent.

It is possible to use the glycidol of formula (V) in racemic or in pure enantiomeric form.

15 Suitable solvents are generally customary organic solvents which do not change under the reaction conditions. These include ethers such as diethyl ether, dioxane or tetrahydrofurane, acetone, dimethylsulfoxide, dimethylformamide or alcohols such as methanol, ethanol, propanol or halogenohydrocarbons such as dichloromethane, trichloromethane or tetrachloromethane. Tetrahydrofurane is preferred.

20


The process is in general carried out in a temperature range from -30°C to +100°C, preferably from 0°C to room temperature.

25 The process is generally carried out at normal pressure. However, it is also possible to carry out it at elevated or reduced pressure (for example in a range from 0.5 to 5 bar).

The compounds of the general formula (III) and (V) are known or can be prepared by customary methods.

5 The compounds of the general formula (IV) are known or as species new and can be prepared characterized in that


[A] first compounds of the general formula (VI)



10 in which

A and R<sup>4</sup> have the abovementioned meaning,

15 are reacted with a catalytic amount of alkali alcoholates, preferred sodium ethanolate to compounds of the general formula (VII)



in which

20 A and R<sup>4</sup> have the abovementioned meaning,

followed by reaction with compounds of the general formula (VIII)




25 in which

R<sup>2</sup> and L have the abovementioned meaning,

in inert solvents, if appropriate in the presence of a base and/or in the presence of an auxiliary to compounds of a general formula (IX)

5



in which


A, L, R<sup>2</sup> und R<sup>4</sup> have the abovementioned meaning,

10

and in a last step the benzyl group is eliminated by hydrogenation,

or


15 [B] compounds of the general formula (X)



in which

20 A, R<sup>1</sup> and R<sup>4</sup> have the abovementioned meaning

are reacted with compounds of the general formula (VIII)



in which

L and R<sup>2</sup> have the abovementioned meaning,


5

in inert solvents, if appropriate in the presence of a base,

and in the case of R<sup>2</sup>/R<sup>3</sup> = H and L = O,

10

compounds of the general formulae (VII) or (X) are reacted with compounds of the general formula (XI)



in which

15

E denotes halogen, preferably chlorine,

and in the case of R<sup>2</sup>/R<sup>3</sup> = H and L = S,

20

compounds of the general formula (X) are reacted with NH<sub>4</sub>SCN,

and in case of R<sup>1</sup>, R<sup>2</sup> and/or R<sup>3</sup> ≠ H the amino groups are derivated optionally by common methods.

25

Suitable solvents for the first step of the procedure [A] (VI-VII) are generally alcohols such as methanol, ethanol or propanol. Ethanol is preferred.

30

Suitable bases for the first step are generally alkali alcoholates such as sodium methanolate, sodium ethanolate or sodium propanolate. Sodium ethanolate is preferred.

The base is employed in catalytic amounts.

The process is in general carried out in a temperature range from 0°C to 60°C, preferably at room temperature.

5

The process is generally carried out at normal pressure. However, it is also possible to carry out it at elevated or reduced pressure (for example in a range from 0.5 to 5 bar).

10

Suitable solvents for the second step of the procedure [A] (VII -> IX) are generally customary organic solvents which do not change under the reaction conditions. These include ethers such as diethyl ether, dioxane or tetrahydrofuran, dimethylsulfoxide, dimethylformamide or halogenohydrocarbons such as dichloromethane, trichloromethane or tetrachloromethane. Dichloromethane is preferred.

15

The process is in general carried out in a temperature range from -30°C to +40°C, preferably from -10°C to room temperature.

The process is generally carried out at normal pressure. However, it is also possible to carry out it at elevated or reduced pressure (for example in a range from 0.5 to 5 bar).

20

The hydroxyl-protective group is in general removed with hydrogen in ethyl acetate, diethyl ether or tetrahydrofuran. Suitable catalysts are noble metal catalysts, preferably palladium and palladium on charcoal .

25

Suitable solvents for the procedure [B] are generally customary organic solvents which do not change under the reaction conditions. These include ethers such as diethyl ether, dioxane or tetrahydrofuran, dimethylsulfoxide, dimethylformamide or halogenohydrocarbons such as dichloromethane, trichloromethane or tetrachloromethane. Dichloromethane is preferred.

30


Suitable bases of the procedure [B], if appropriate, are generally inorganic or organic bases. These preferably include alkali metal hydroxides such as, for example, sodium hydroxide, sodium hydrogencarbonate or potassium hydroxide, alkaline earth metal hydroxides such as, for example, barium hydroxide, alkali metal carbonates such as calcium carbonate, or alkaline metal or organic amines (trialkyl(C<sub>1</sub>-C<sub>6</sub>)amines) such as triethylamine, or heterocycles such as 1,4-diazabicyclo[2.2.2]octane (DABCO), 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), or amides such as sodium amides, lithium butyl amide or butyllithium, pyridine or methylpiperidine. It is also possible to employ alkali metals, such as sodium or its hydrides such as sodium hydride, as bases. Potassium carbonate, triethylamine, sodium hydrogencarbonate and sodium hydroxide are preferred.

The base is employed in an amount from 1 mol to 10 mol, preferably from 1.0 mol to 4 mol, relative to 1 mol of the compounds of the general formulae (VIII).

The process is in general carried out in a temperature range from -30°C to +100°C, preferably from -10°C to +50°C.

The process is generally carried out at normal pressure. However, it is also possible to carry out it at elevated or reduced pressure (for example in a range from 0.5 to 5 bar).


The compounds of the general formula (VI) are as species new and can be prepared by reaction of compounds of the general formula (XII)



in which

A has the abovementioned meaning,

with hydroxylamine hydrochloride in a presence of sodium formate to compounds of the general formula (XIII)



in which

A has the abovementioned meaning,

10 followed by reaction with compounds of the general formula (XIV)



in which

15

R<sup>4</sup> has the abovementioned meaning,

and

20 T represents halogen, preferably bromine,

in inert solvents and in the presence of a base.

25 Suitable solvents are generally customary organic solvents which do not change under the reaction conditions. These include ethers such as diethyl ether, dioxane or tetrahydrofurane, acetone, dimethylsulfoxide, dimethylformamide or alcohols such as methanol, ethanol, propanol or halogenohydrocarbons such as dichloromethane, trichloromethane or tetrachloromethane. Acetone and dimethylformamide are preferred.

Suitable bases for the procedure are generally inorganic or organic bases. These preferably include alkali metal hydroxides such as, for example, sodium hydroxide or potassium hydroxide, alkaline earth metal hydroxides such as, for example, barium hydroxide, alkali metal carbonates such as sodium carbonate, potassium carbonate,  
5 alkaline earth metal carbonates such as calcium carbonate. Potassium carbonate (powdered) is preferred.

The base is employed in an amount from 1 mol to 10 mol, preferably from 1.0 mol to  
10 4 mol, relative to 1 mol of the compounds of the general formulae (XIII).

The process is in general carried out in a temperature range from -30°C to +100°C, preferably from -10°C to +60°C.

15 The process is generally carried out at normal pressure. However, it is also possible to carry out it at elevated pressure or at reduced pressure (for example in a range from 0.5 to 5 bar).

20 The compounds of the general formula (XII) are known or as species new and can be prepared by reaction of 2,4-dihydroxy-benzaldehydes with benzylbromide in one of the abovementioned solvents and bases, preferably in acetone with potassium carbonate at room temperature.

25 The compounds of the general formula (XIII) are new and can be prepared like described above.

The compounds of the general formula (XIV) are known or as species new and can be prepared by customary methods.

30 The compounds of the general formulae (III), (V), (VIII), (IX) and (X) are known or as species new and can be prepared by customary methods.



5 The compounds of the general formula (VII) can be prepared like described above or in a single step procedure by reacting compounds of the general formula (XIII) with compounds of the general formula (XIV) in the presence of a surplus of sodium ethylate under reflux .

10 Surprisingly it was found that compounds given by the general formula (I) inhibited oxygen radical formation as well as  $\text{TNF}\alpha$  (tumor necrosis factor) production, but potentiated the release of IL-10. These compounds elevated cellular cyclic AMP probably by inhibition of phagocyte phosphodiesterase activity.

15 The compounds according to the invention specifically inhibit the production of superoxide by polymorphonuclear leukocytes (PMN). Furthermore, these compounds inhibit  $\text{TNF}\alpha$  release and potentiate IL-10 production in human monocytes in response to a variety of stimuli including bacterial lipopolysaccharide (LPS), complement-opsonized zymosan (ZymC3b) and IL-1 $\beta$ .

20 The described effects are probably mediated by the elevation of cellular cAMP probably due to inhibition of the type IV phosphodiesterase responsible for its degradation.

They can therefore be employed in medicaments for the treatment of acute and chronic inflammatory processes.

25 The compounds according to the invention are preferably suitable for the treatment and prevention of acute and chronic inflammation and auto immune diseases, such as emphysema, alveolitis, shock lung, all kinds of asthma, COPD, ARDS, bronchitis, arteriosclerosis, arthrosis, inflammations of the gastro-intestinal tract, rheumatoid arthritis, myocarditis, sepsis and septic shock, arthritis, rheumatoid spondylitis and  
30 osteoarthritis, gram negative sepsis, toxic shock syndrome, acute respiratory distress syndrome, bone resorption diseases, reperfusion injury, graft vs host reaction, allograft

rejection, malaria, myalgias, HIV, AIDS, cachexia, Cronh's disease, ulcerative colitis, pyresis, system lupus erythematosus, multiple sclerosis, type I diabetes mellitus, psoriasis, Bechet's disease, anaphylactoid purpura nephritis, chronic glomerulo-nephritis, inflammatory bowel disease and leukemia. The compounds according to the invention are additionally suitable for reducing the damage to infarct tissue after reoxygenation. In this case the simultaneous administration of allopurinol to inhibit xanthine oxidase is of advantage. Combination therapy with superoxide dismutase is also of use.

#### 10 Test description

##### 1. Preparation of human PMN

Blood was taken from healthy subjects by venous puncture and neutrophils were purified by dextran sedimentation and resuspended in the buffered medium.

##### 2. Inhibition of FMLP-stimulated production of superoxide radical anions.

Neutrophils ( $2.5 \times 10^5 \text{ ml}^{-1}$ ) were mixed with cytochrome C (1.2 mg/ml) in the wells of a microtitre plate. Compounds according to the invention were added in dimethyl sulphoxide (DMSO). Compound concentration ranged from 2.5 nM to 10  $\mu\text{M}$ , the DMSO concentration was 0.1% v/v in all wells. After addition of cytochalasin b ( $5 \mu\text{g} \times \text{ml}^{-1}$ ) the plate was incubated for 5 min at 37°C. Neutrophils were then stimulated by addition of  $4 \times 10^{-8} \text{ M}$  FMLP and superoxide generation measured as superoxide dismutase inhibitable reduction of cytochrome C by monitoring the  $\text{OD}_{550}$  in a microtitre plate spectrophotometer, such as a Thermomax microtitre plate spectrometer. Initial rates were calculated using a kinetic calculation program, e.g. a softmax programme. Blank wells contained 200 units of superoxide dismutase.

30 The inhibition of superoxide production was calculated as follows:

$$\frac{[1 - ((R_x - R_b))]}{(R_o - R_b)} \cdot 100 = \% \text{ inhibition}$$

R<sub>x</sub> = Rate of the well containing the compound according to the invention.

R<sub>o</sub> = Rate in the control well.

5 R<sub>b</sub> = Rate in the superoxide dismutase containing blank well.

Compounds according to the invention have IC<sub>50</sub> values in the range 0,001 µM-1µM. Example 6 exhibits a IC<sub>50</sub>(O<sub>2</sub><sup>-</sup>)-value of 0.08µM.

10 3. Measurement of PMN cyclic AMP concentration

The compounds according to the invention were incubated with 3.7 x 10<sup>6</sup> PMN for 5 min at 37°C before addition of 4 x 10<sup>-8</sup> M FMLP. After 6 min protein was precipitated by the addition of 1% v/v conc. HCl in 96% v/v ethanol containing 0.1 mM EDTA. After centrifugation the ethanolic extracts were evaporated to dryness under N<sub>2</sub> and resuspended in 50 mM Tris/HCl pH 7.4 containing 4 mM EDTA. The cyclic AMP concentration in the extracts was determined using a cyclic AMP binding protein assay supplied by Amersham International plc. Cyclic AMP concentrations were expressed as percentage of vehicle containing control incubations.

20 Compounds elavate the cAMP-level at 1 µM compound 0-400% of control values.

4. Assay of PMN phosphodiesterase

25 This was performed as a particulate fraction from human PMN essentially as described by Souness and Scott (Biochem. J. 291, 389-395, 1993). Particulate fractions were treated with sodium vanadate / glutathione as described by the authors to express the descrete stereospecific site on the phosphodiesterase enzyme. Compounds according to the invention had IC<sub>50</sub> values ranging from 0.001 µM to 10 µM. Example 6 exhibits a IC<sub>50</sub>(PDE IV) of 0.17 µM.

30

5. Assay of human platelet phosphodiesterase

5 This was performed essentially as described by Schmidt et al. (Biochem. Pharmacol. 42, 153-162, 1991) except that the homogenate was treated with vanadate glutathione as above. Compounds according to the invention had IC<sub>50</sub> values greater than 100 µM.

6. Assay of binding to the rolipram binding site in rat brain membranes

10 This was performed essentially as described by Schneider et al. (Eur. J. Pharmacol. 127, 105-115, 1986). Compounds according to the invention had IC<sub>50</sub> values in the range 0,01 to 10 µM.

7. Preparation of human monocytes

15 Blood was taken from normal donors. Monocytes were isolated from peripheral blood by density centrifugation, followed by centrifugal elutriation.

8. Endotoxin induced TNF release

20 Monocytes (1 x 10<sup>6</sup> ml<sup>-1</sup>) were stimulated with LPS (2 µg ml<sup>-1</sup>) and coincubated with the compounds at different concentrations (10<sup>-4</sup> to 10 µg ml<sup>-1</sup>). Compounds were dissolved in DMSO/medium (2% v/v). The cells were incubated in RPMI-1640 medium glutamine/FCS supplemented and at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>. After 18 to 24 hours TNF was determined in the supernatants by an human TNF specific ELISA (medgenix). Controls were nonstimulated and LPS stimulated monocytes without compounds.

25 9. Endotoxin induced shock lethality in mice

30 B6D2F1 mice (n=10) were sensitized with galactosamine (600 mg/kg), and shock and lethality were triggered by LPS (0.01 µg/mouse). The compounds were administered intravenously 1 hour prior LPS. Controls were LPS challenged mice without compound. Mice were dying 8 to 24 hours post LPS challenge.

The galactosamine / LPS mediated mortality was reduced.

10. Stimulation of human monocytes and determination of cytokine levels

5 Human monocytes ( $2 \times 10^5$  in 1 ml) were stimulated with 100 ng/ml LPS, 0.8 mg/ml zymC3b or 10 ng/ml IL-1 $\beta$  in the presence of test compounds. The final DMSO concentration was maintained at 0.1 % v/v. Cells were incubated overnight in a humidified atmosphere of 5% CO<sub>2</sub> at 37°C. Supernatants were harvested and stored at -70°C. The TNF $\alpha$  concentration was measured by  
10 ELISA using the A6 anti-TNF monoclonal antibody (Miles) as the primary antibody. The secondary antibody was the polyclonal anti-TNF $\alpha$  antibody IP300 (Genzyme) and the detection antibody was a polyclonal anti-rabbit IgG alkaline phosphatase conjugate (Sigma). IL-10 was determined by ELISA (Biosource).

15

The new active compounds can be converted in a known manner into the customary formulations, such as tablets, coated tablets, pills, granules, aerosols, syrups, emulsions, suspensions and solutions, using inert, nontoxic, pharmaceutically suitable excipients or solvents. In this connection, the therapeutically active compound should in each case  
20 be present in a concentration of about 0.5 to 90% by weight of the total mixture, i.e. in amounts which are sufficient in order to achieve the dosage range indicated.

The formulations are prepared, for example, by extending the active compounds with solvents and/or excipients, if appropriate using emulsifiers and/or dispersants, where,  
25 for example, in the case of the use of water as a diluent, organic solvents can be used as auxiliary solvents if appropriate.

Administration is carried out in a customary manner, preferably orally or parenterally.

30 In the case of parenteral administration, solutions of the active compound can be employed using suitable liquid vehicles.

In general, it has proved advantageous on intravenous administration to administer amounts from about 0.001 to 10 mg/kg, preferably about 0.01 to 5 mg/kg of body weight to achieve effective results, and on oral administration the dosage is about 0.01 to 25 mg/kg, preferably 0.1 to 10 mg/kg of body weight.

In spite of this, it may be necessary to depart from the amounts mentioned, in particular depending on the body weight or the type of application route, on individual behaviour towards the medicament, the manner of its formulation and the time or interval at which administration takes place. Thus, in some cases it may be sufficient to manage with less than the abovementioned minimum amount, while in other cases the upper limit mentioned must be exceeded. In the case of administration of relatively large amounts, it is advisable to divide these into several individual doses over the course of the day.


15

**Solvents**

	a	=	dichloromethane/methanol	15:1 (v/v)
	b	=	dichloromethane/methanol	20:1
20	c	=	dichloromethane/methanol	50:1
	d	=	dichloromethane/methanol	40:1
	e	=	ethylacetate/cyclohexane	5:1
	f	=	chloroform/methanol	10:1
	g	=	dichloromethane/methanol	10:1
25	h	=	dichloromethane/methanol	30:1
	i	=	chloroform/methanol	30:1
	j	=	chloroform/methanol/water/acetic acid	70:30:5:5
	k	=	dichloromethane/methanol/formic acid	9:1:0.1
	l	=	dichloromethane/methanol/NH <sub>3</sub>	9:1:0.1
30	m	=	dichloromethane/methanol	5:1
	n	=	dichloromethane/methanol	7:1
	o	=	dichloromethane/methanol/NH <sub>3</sub>	12:1:0.1

**Starting compounds:****Example I**

5

**4-Benzyloxy-2-hydroxybenzaldehyde**


10 13.8 g (0.1 mol) 2,4-dihydroxybenzaldehyde are dissolved in 150 ml acetone, 17.1 g (0.1 mol) benzylbromide and 13.8 g (0.1 mol) potassium carbonate are added, and the mixture is stirred at room temperature for 3 d. After filtration, the solvent is removed in vacuo and the residual crude product further purified by column chromatography on silica gel using dichloromethane as eluent.

15 Yield: 9.2 g (40%)

m.p.: 81-82°C

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz): δ = 5.11 (s, 2H; CH<sub>2</sub>); 6.51 (d, 1H, Ar-H); 6.61 (dd, 1H, Ar-H); 7.32 - 7.45 (m, 6H, Ar-H); 9.82 (s, 1H, CHO); 11.60 (s, 1H, OH) ppm.

20 **Example II**

**4-Benzyloxy-2-hydroxybenzonitrile**

25

25.0 g (0.11 mol) of the example I, 8.67 g (0.13 mol) hydroxylamine hydrochloride and 15.0 g (0.22 mol) sodium formate are dissolved in 170 ml formic acid, and the solution

stirred under reflux for 1-3 h. The reaction mixture is cooled to room temperature, poured into ice/water and extracted with dichloromethane. The organic layer is dried ( $\text{Na}_2\text{SO}_4$ ), the solvent removed in vacuo and the residue triturated with dichloromethane. Alternatively, the precipitate which is formed after dilution with ice/water is filtered off and dried in a desiccator in the presence of phosphorus pentoxide. The crude product is crystallized from dichloromethane/cyclohexane.


Yield: 15.0 g (61%)

m.p.: 135-136°C

$^1\text{H-NMR}$  (400 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  = 5.13 (s, 2H,  $\text{CH}_2$ ); 6.58 (d, 1H, Ar-H); 6.60 (dd, 1H, Ar-H); 7.32 - 7.46 (m, 5H, Ar-H); 7.50 (d, 1H, Ar-H); 11.10 (br.s, 1H, OH) ppm.

### Example III

$\omega$ -Bromo-2,4-dichloroacetophenone



To a solution of 155.2 g (788 mmol) 2,4-dichloroacetophenone in 450 ml glacial acetic acid are added 40.4 ml (788 mmol) bromine at 50°C. After 1 h at 50°C, further 4 ml (78.8 mmol) bromine are added and the mixture is stirred for an additional hour. The solution is cooled to room temperature, concentrated to half of its volume and diluted with 800 ml water. The mixture is neutralized with solid sodium carbonate and extracted three times with ethyl acetate. The organic layer is dried ( $\text{Na}_2\text{SO}_4$ ), the solvent removed in vacuo, finally under high vacuum. The crude product is used directly without further purification for the following reaction and stored in the meantime at -20°C.


Yield: 211.4 g (quant.); yellow oil



**Example IV**

3-Amino-2-(2,4-dichlorobenzoyl)-6-benzyloxy-benzofuran

5



a) Single step procedure:

10.0 g (44.0 mmol) of example II are dissolved in 200 ml ethanol, and 5.98 g (88.0 mmol) sodium ethylate and 15.4 g (48.0 mmol) of example III are added. The mixture is stirred under reflux for 4 d. During this time further 12.0 g sodium ethylate and 7.7 g of example III are added. The solvent is removed in vacuo and the residue is purified by column chromatography on silica gel (eluent: 1. dichloromethane, 2. dichloromethane/methanol 98.2).

15

Yield: 8.1 g (44%)

m.p.: 158°C

NMR-data see below.


**Example V**

20

Two step procedure:

a)  $\omega$ -[(5-Benzyloxy-2-cyan)-phenoxy]-2,4-dichloroacetophenone:

- 32 -



To a mixture of 26.6 g (118 mmol) of example II and 48.85 g (354 mmol) powdered potassium carbonate in 800 ml acetone are added 34.8 g (130 ml) of example III under reflux. After a further 1-2 h under reflux, the mixture is filtered, the solvent evaporated in vacuo and the residue triturated using dichloromethane (1st crop of product) and dichloromethane/cyclohexane (4:1) (2nd crop of product).

Yield: 26.2 g (54%)

b) Two step procedure:

To a solution of 37.5 g (91 mmol) of the compound of example Va in 550 ml ethanol, three pellets of sodium hydroxide are added and the mixture is stirred at 50°C for 30 min. After cooling to room temperature, the mixture is filtered off (1st crop of product) and the filtrate is diluted with 300 ml of water. Most of the ethanol is removed in vacuo and the formed precipitate is filtered (2nd crop of product).


Yield: 36.7 g (98%)

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 400 MHz): δ = 5.14 (s, 2H, CH<sub>2</sub>); 6.97 (dd, 1H, Ar-H); 7.10 (d, 1H, Ar-H); 7.31 - 7.47 (m, 5H, Ar-H); 7.53 (br.s, 2H, NH<sub>2</sub>); 7.57 (s, 2H, Ar-H); 7.75 (s, 1H, Ar-H); 7.93 (d, 1H, Ar-H); ppm.

Ms (EI): 411 (M<sup>+</sup>)

**Example VI**

N-[2-(2,4-Dichlorobenzoyl)-6-benzyloxy-benzofuran-3-yl]urea



5

30.9 g (75 mmol) of example IV are dissolved in 750 ml dichloromethane and cooled to 0°C. 11.6 g (82 mmol) chlorosulfonyl isocyanate in dichloromethane are added and the mixture is stirred at 0°C for 30 min and at room temperature for 3 h. The solvent is removed in vacuo, the residue suspended in water and stirred vigorously for 3 h at 60°C. After cooling to r.t. the suspension is filtered, the crude product dried in a desiccator in the presence of phosphorus pentoxide and triturated with diisopropylether at 40°C.

10

Yield: 33.7 g (98%)

m.p.: 192-193°C

15


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 400 MHz): δ = 5.18 (s, 2H, CH<sub>2</sub>); 7.00 (dd, 1H, Ar-H); 7.05 (br.s, 2H, NH<sub>2</sub>); 7.20 (d, 1H, Ar-H); 7.30 - 7.47 (m, 5H, Ar-H); 7.61 (dd, 1H, Ar-H); 7.68 (d, 1H, Ar-H); 7.82 (d, 1H, Ar-H); 8.37 (d, 1H, Ar-H); 9.71 (s, 1H, NH) ppm.

MS (FAB): 455 (M+H)<sup>+</sup>

20

**Example VII**

N-[2-(2,4-Dichlorobenzoyl)-6-hydroxy-benzofuran-3-yl]urea



5

To a solution of 5.0 g of example VI in 100 ml THF are added 250 mg 10% palladium on activated charcoal, and the mixture is hydrogenated at atmospheric pressure and room temperature for 1-3 d. If necessary, further 125 mg 10% Pd/C are added and the hydrogenation is continued for 24 h. The mixture is filtered through celite and the solvent is removed in vacuo to about 1/3 of the original volume. Dichloromethane is added, the mixture is stirred at 0°C and the formed precipitate is filtered off.

10

Yield: 4.0 g (quant.)

m.p.: 230-231°C

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 400 MHz): δ = 6.73 (d, 1H, Ar-H); 6.80 (dd, 1H, Ar-H); 7.03 (br.s, 2H, NH<sub>2</sub>); 7.60 (dd, 1H, Ar-H); 7.65 (d, 1H, Ar-H); 7.80 (s, 1H, Ar-H); 8.30 (d, 1H, Ar-H); 9.72 (s, 1H, NH); 10.40 (br.s, 1H, OH) ppm.


15

MS (EI): 364 (M<sup>+</sup>).

In analogy to this procedure the following examples shown in table B are prepared:

20


Table B:



Ex.-No.	R <sup>14</sup>	R <sup>15</sup>	R <sup>16</sup>	R <sup>17</sup>	Yield(% of th.)	R <sub>f</sub> *
VIII	F	H	F	H	90	0,76 <sup>b)</sup>
IX	H	H	Cl	H	40	0,38 <sup>a)</sup>
X	OCH <sub>3</sub>	H	OCH <sub>3</sub>	H	90	0,38 <sup>b)</sup>
XI	Cl	H	H	H	91	0,36 <sup>a)</sup>
XII	Cl	H	H	Cl	91	0,31 <sup>b)</sup>
XIII	CH <sub>3</sub>	H	CH <sub>3</sub>	H	58	0,26 <sup>b)</sup>

5

## Example XVI



10 15 g (41.1 mmol) 6-hydroxybenzofurane, 2.82 ml (41.1 mmol) R-glycidol, and 10.8 g (41.1 mmol) triphenylphosphin are dissolved in 200 ml dry THF. At 0°C 6.43 ml (41.1 mmol) diethyl azodicarboxylate (DEAD) are added dropwise, and the reaction mixture is stirred at room temperature for a total of 4 days. During this time, three

additional portions of glycidol (0.94 ml, 13.7 mmol each), triphenylphosphin (3.6 g, 13.7 mmol each), and DEAD (2.1 ml, 13.7 mmol each) are added to the reaction mixture. The mixture is filtered (first crop of product), the filtrate concentrated to half of its original volume and filtered again (second crop of product). The filtrate is diluted  
5 with dichloromethane, washed with water twice, dried over sodium sulfate, and concentrated to a volume of about 200 ml. An equal volume of diethylether is added, the mixture stirred for 15 min, and filtered again to yield a third crop of product.

Combined yield: 15.89 g (91.8% of th.)

$R_f = 0.57$  (chloroform/methanol (15:1))





10






**Preparation examples:**

**General procedure for the synthesis of benzofuranylamino alcohols:**

15 The benzofuranyl epoxide is dissolved in 1:1 dry tetrahydrofurane/methanol, 2 mol-equivalent of the amine component are added, and the reaction mixture is stirred at 40 - 50°C for 1-7 days. After concentration, the residue is taken up in dichloromethane or chloroform, and washed with water several times. The organic layer is dried over sodium sulfate, concentrated, and the crude product purified by column chromatography over silica gel.  
20

The following examples were prepared in analogy to the general procedure:

Ex.- No.	Structure	Yield (%) of theory)	R <sub>f</sub> *
1		17.1	0.15 (g)
2		52.1	0.29 (l)
3		22.1	0.31 (m)
4		21.8	0.46 (m)


Ex.- No.	Structure	Yield (% of theory)	R <sub>f</sub> *
5		46.5	0.28 (n)
6		31.5	0.40(g)
7		61.2	0.40(g)
8		14.3	0.33 <sup>o)</sup>
9		31.0	0.25 <sup>o)</sup>



We claim:

1. Benzofuranylaminoholols of the general formula (I)

5



in which

10 A represents hydrogen, straight-chain or branched acyl or alkoxycarbonyl, each having up to 6 carbon atoms, halogen, carboxyl, cyano, nitro, hydroxyl, trifluoromethyl or trifluoromethoxy, or straight-chain or branched alkyl having 1 to 6 carbon atoms, which is optionally substituted by carboxyl, alkoxy or alkoxycarbonyl each having 1 to 4 carbon atoms, phenoxy or benzoyl,

15

R<sup>1</sup> represents hydrogen, straight-chain or branched alkyl having 1 to 4 carbon atoms, an amino protecting group or a group of the formula -CO-R<sup>7</sup>

20

in which

R<sup>7</sup> denotes straight chain or branched alkoxy having 1 to 4 carbon atoms,

25

R<sup>2</sup> and R<sup>3</sup> are identical or different and represent hydrogen, cycloalkyl having 3 to 6 carbon atoms, straight chain or branched alkyl, alkoxycarbonyl or alkenyl each having 1 to 8 carbon atoms,

or

R<sup>2</sup> and R<sup>3</sup> together with the nitrogen atom form a 5- to 7-membered saturated heterocycle optionally having a further oxygen atom,

5

R<sup>4</sup> represents aryl having 6 to 10 carbon atoms or represents a 5- to 7-membered, aromatic, saturated or unsaturated heterocycle, which can contain 1 to 3 oxygen, sulphur and/or nitrogen atoms as heteroatoms or a residue of a formula -NR<sup>8</sup>,

10

wherein

R<sup>8</sup> denotes hydrogen or straight-chain or branched alkyl or alkoxy-carbonyl each having 1 to 6 carbon atoms,

15

and to which further a benzene ring can be fused and wherein aryl and/or the heterocycle are optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising hydroxyl, halogen, nitro, 1H-tetrazolyl, pyridyl, trifluoromethyl, trifluoromethoxy, difluoromethyl, difluoromethoxy, cyano, carboxy, straight-chain or branched alkoxy, alkoxy-carbonyl or acyl each having 1 to 6 carbon atoms or by straight-chain or branched alkyl having 1 to 5 carbon atoms, which is optionally substituted by carboxyl or straight-chain or branched alkoxy-carbonyl having 1 to 4 carbon atoms or by a group of formula -NR<sup>9</sup>R<sup>10</sup>, -SR<sup>11</sup>, -(NH)<sub>a</sub>-SO<sub>2</sub>R<sup>12</sup> or -O-SO<sub>2</sub>R<sup>13</sup>,

20

25

in which

R<sup>9</sup> and R<sup>10</sup> are identical or different and denote hydrogen or a straight-chain or branched alkyl having 1 to 4 carbon atoms,

30

or

$R^9$  denotes hydrogen

5

and

$R^{10}$  denotes straight-chain or branched acyl having 1 to 6 carbon atoms,

10

$R^{11}$  denotes hydrogen or straight-chain or branched alkyl having 1 to 4 carbon atoms,

$a$  denotes a number 0 or 1,

15

$R^{12}$  and  $R^{13}$  are identical or different and represent straight-chain or branched alkyl having 1 to 6 carbon atoms, benzyl or phenyl, which are optionally substituted by trifluoromethyl, halogen or straight-chain or branched alkyl having 1 to 4 carbon atoms,

20

$L$  represents an oxygen or sulfur atom

25

$R^5$  and  $R^6$  represents hydrogen or straight-chain or branched alkyl having 1 to 6 carbon atoms, which is optionally substituted by aryl having 6 to 10 carbon atoms or by a 5- to 7-membered aromatic, saturated or unsaturated heterocycle having 1 to 3 heteroatoms from the series comprising N, S, O and/or a residue of a formula  $-NR^{14}$

wherein

30

$R^{14}$  denotes hydrogen or straight-chain or branched alkyl or alkoxy-carbonyl each having 1 to 6 carbon atoms

5 and to which a phenyl ring can be fused and which are optionally monosubstituted to disubstituted by identical or different substituents from the series comprising halogen, cyano or by a straight-chain or branched alkyl or alkoxycarbonyl each having 1 to 6 carbon atoms,

or represent a 6-membered saturated N-heterocycle, which is optionally substituted by alkoxycarbonyl having 1 to 6 carbon atoms,

10 or

R<sup>5</sup> and R<sup>6</sup> together with the nitrogen atom form a 5- to 6-membered, aromatic, saturated or unsaturated heterocycle having 1 to 3 heteroatoms from the series comprising N, S, O and/or a residue of a formula -NR<sup>14</sup>,

15

and to which a phenyl ring can be fused and which is optionally monosubstituted to disubstituted by identical or different substituents from the series comprising halogen, cyano or by a straight-chain or branched alkyl or alkoxycarbonyl each having 1 to 6 carbon atoms,

20

and salts thereof.

2. Compounds according to claim 1, in which

25 A represents hydrogen, halogen, carboxyl, cyano, nitro, hydroxyl, trifluoromethyl, trifluoromethoxy or alkoxy having 1 to 4 carbon atoms

R<sup>1</sup> represents hydrogen, straight-chain or branched alkyl having 1 to 4 carbon atoms or a group of the formula -CO-R<sup>7</sup>

30

in which

$R^7$  denotes straight chain or branched alkoxy having 1 to 4 carbon atoms,

5  $R^2$  and  $R^3$  are identical or different and represent hydrogen, cyclobutyl, cyclopentyl, cyclohexyl or straight-chain or branched alkyl, alkoxy carbonyl or alkenyl each having 1 to 4 carbon atoms,

or

10

$R^2$  and  $R^3$  together with the nitrogen atom form a pyrrolidinyl-, piperidinyl- or morpholinyl-ring,

and

15

$R^4$  represents phenyl, pyridyl or thienyl, wherein all rings are optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising hydroxyl, fluorine, chlorine, bromine, nitro, carboxy, straight-chain or branched alkoxy, alkoxy carbonyl or acyl each having 1 to 3 carbon atoms, or by methyl, ethyl, n-propyl or isopropyl which is optionally substituted by carboxyl or straight-chain or branched alkoxy carbonyl having up to 3 carbon atoms

20

L represents an oxygen or sulfur atom,

25

$R^5$  and  $R^6$  represents hydrogen or straight-chain or branched alkyl having 1 to 4 carbon atoms, which is optionally substituted by phenyl, pyridyl, imidazolyl, pyrrol, morpholinyl, piperidinyl, piperazinyl, pyrrolidinyl, wherein the heterocycles optionally contain a residue of a formula  $-NR^{14}$ ,

30

wherein

$R^{14}$  denotes hydrogen or straight-chain or branched alkyl or alkoxy-carbonyl each having 1 to 4 carbon atoms,

and wherein the ring systems are optionally monosubstituted by straight-chain or branched alkyl, alkoxy or alkoxycarbonyl each having 1 to 3 carbon atoms,

or

$R^5$  and  $R^6$  together with the nitrogen atom form a pyrazolyl-, triazolyl-, tetrazolyl-, imidazolyl-, pyrrol-, morpholinyl-, piperidinyl-, pyrrolidinyl-, piperazinylring, wherein the heterocycles optionally contain a residue of a formula  $-NR^{14}$ ,

and wherein the ringsystem is optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising halogen, cyano or by a straight-chain or branched alkyl having 1 to 6 carbon atoms,

and salts thereof.

3. Compounds according to claim 1 or 2, in which

A represents hydrogen,

$R^1$  represents hydrogen, methyl, ethyl, n-propyl or isopropyl or a group of the formula  $-CO-R^7$ ,

in which

R<sup>7</sup> denotes methoxy, ethoxy, n-propoxy or isopropoxy,

R<sup>2</sup> and R<sup>3</sup> represent hydrogen,

5

R<sup>4</sup> represents phenyl or pyridyl, which are optionally up to difold substituted by identical or different substituents from the series fluorine, chlorine, methyl or methoxy,

10

L represents an oxygen atom,

15

R<sup>5</sup> and R<sup>6</sup> represents hydrogen or straight-chain or branched alkyl having 1 to 4 carbon atoms, which is optionally substituted by phenyl, pyridyl, imidazolyl, pyrrol, piperidiny, wherein the heterocycles optionally contain a residue of a formula -NR<sup>14</sup>,

wherein

20

R<sup>14</sup> denotes hydrogen or straight-chain or branched alkyl or alkoxy-carbonyl each having 1 to 3 carbon atoms,

25

and wherein the ring systems are optionally monosubstituted by straight-chain or branched alkyl, alkoxy or alkoxycarbonyl each having 1 to 3 carbon atoms,

or represent a 6-membered saturated N-heterocycle, which is optionally substituted by alkoxycarbonyl having 1 to 6 carbon atoms,

or



30

R<sup>5</sup> and R<sup>6</sup> together with the nitrogen atom form a imidazolyl-, pyrrol-, morpholinyl-, piperidinyl-, pyrrolidinyl-, piperazinylring, wherein the heterocycles are optionally contain a residue of a formula -NR<sup>14</sup>,




5 and wherein the ringsystem are optionally monosubstituted to trisubstituted by identical or different substituents from the series comprising by a straight-chain or branched alkyl or alkoxycarbonyl each having 1 to 3 carbon atoms,





10 and salts thereof.

4. Compounds according to any one of claims 1 to 3, selected from the group consisting of:










5. A process for the preparation of compounds according to claim 1, characterized in that compounds of the general formula II



in which

A, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and L have the abovementioned meaning,

are reacted with amines of the general formula (III)



in which

R<sup>5</sup> and R<sup>6</sup> have the abovementioned meaning,

in the presence of an inert solvent.

6. Benzofuranylaminoalcohols according to any one of claims 1 to 4 for therapeutic use.
7. The composition containing at least one Benzofuranylaminoalcohol according to any one of claims 1 to 4 and a pharmacologically acceptable diluent.
8. A composition according to claim 7 for the treatment and prevention of acute and chronic inflammatory processes.

- 5
9. The process for the preparation of composition according to claim 7 and 8 characterized in that the Benzofuranylaminoalcohol together with customary auxiliaries is brought into a suitable application form.
10. Use of Benzofuranylaminoalcohols according to anyone of claims 1 to 4 for the preparation of medicaments.
- 10 11. Use according to claim 10 for the preparation of medicaments for the treatment and prevention of acute and chronic inflammatory processes.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 November 2000 (23.11.2000)

PCT

(10) International Publication Number  
**WO 00/69841 A3**

(51) International Patent Classification<sup>7</sup>: **C07D 307/83**,  
307/81, 405/12, A61P 29/00, A61K 31/44, 31/343 //  
(C07D 405/12, 307:00, 213:00)

(21) International Application Number: **PCT/EP00/04015**

(22) International Filing Date: **4 May 2000 (04.05.2000)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
9911453.0 **17 May 1999 (17.05.1999) GB**

(71) Applicant (*for all designated States except US*): **BAYER AKTIENGESSELLSCHAFT [DE/DE]**; D-51368 Leverkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BRÄUNLICH, Gabriele [DE/DE]**; Claudiusweg 9, D-42115 Wuppertal (DE). **ES-SAYED, Mazen [DE/DE]**; Ricarda-Huch-Str. 36, D-40764 Langenfeld (DE). **FISCHER, Rüdiger [DE/DE]**; Zu den Fussfällen 23, D-50529 Pulheim (DE). **FUGMANN, Burkhard [DE/DE]**; Dr.-Kessel-Strasse 3, D-40878 Ratingen (DE). **HENNING, Rolf [DE/DE]**; Am Jagdhaus 115, D-42113 Wuppertal (DE). **SCHNEIDER, Stephan [DE/DE]**; Am Ringofen 46, D-42327 Wuppertal (DE). **SPERZEL, Michael [DE/DE]**; Normanenstrasse 31, D-42275 Wuppertal (DE). **SCHLEMMER, Karl-Heinz [DE/DE]**; Wildsteig 22a, D-42113 Wuppertal (DE). **STURTON, Graham [GB/GB]**; 184 Windsor Road, Bray Maidenhead SL6 2DW (GB). **FITZGERALD, Mary [GB/GB]**; 2 Paternoster Court, Cassington

Road Yarton, Oxford OX5 1QB (GB). **BRIGGS, Barbara [GB/GB]**; 24 Latchmere Road, Kingston-upon-Thames, Surrey KT2 5TW (GB). **CONCEPTION, Arnel [PH/JP]**; 3-6-27-207, Shihatsuji-cho, Nara City, Nara 630-8114 (JP). **BULLOCK, William [US/US]**; 60 Hermann Lane, Easton, CT 06612 (US).

(74) Common Representative: **BAYER AKTIENGESSELLSCHAFT**; D-51368 Leverkusen (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


— *with international search report*

(88) Date of publication of the international search report:  
**2 May 2002**

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 00/69841 A3

(54) Title: **BENZOFURANYLAMINOALCOHOLS**



(0)

(57) Abstract: Benzofuranyl aminoalcohols of general formula (I), process for their preparation and their use in medicaments, especially for the treatment of inflammatory processes.

# INTERNATIONAL SEARCH REPORT

national Application No  
PCT/EP 00/04015

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D307/83 C07D307/81 C07D405/12 A61P29/00 A61K31/44  
A61K31/343 //(C07D405/12,307:00,213:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61P A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EP0-Internal, PAJ, WPI Data, CHEM ABS Data, BEILSTEIN Data, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 731 099 A (BAYER AG) 11 September 1996 (1996-09-11) cited in the application claim 1; examples 2,56; tables 1-6 ---	1-11
Y	EP 0 779 291 A (BAYER AG) 18 June 1997 (1997-06-18) claim 1; table 1 ---	1-11
Y	WO 98 02440 A (BRAEUNLICH GABRIELE ;SCHNEIDER STEPHAN (DE); BAYER AG (DE); ES SAY) 22 January 1998 (1998-01-22) claim 1; examples 17,18,37 ---	1-11
Y	EP 0 146 243 A (MERCK FROSST CANADA INC) 26 June 1985 (1985-06-26) claim 1 -----	1-4

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

7 August 2000

Date of mailing of the international search report

17/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Härtinger, S

# INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/EP 00/04015

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0731099	A	11-09-1996	AT 193704 T	15-06-2000
			AU 705667 B	27-05-1999
			AU 4582896 A	19-09-1996
			BG 100396 A	31-12-1996
			BR 9600930 A	30-12-1997
			CA 2170824 A	07-09-1996
			CN 1134419 A	30-10-1996
			CZ 9600667 A	11-09-1996
			DE 69608721 D	13-07-2000
			FI 960998 A	07-09-1996
			HR 960071 A	31-12-1997
			HU 9600549 A	28-07-1997
			JP 8253466 A	01-10-1996
			NO 960884 A	09-09-1996
			NZ 286107 A	26-08-1998
			PL 313070 A	16-09-1996
			SK 29796 A	01-10-1996
			TR 960863 A	21-10-1996
			ZA 9601774 A	17-09-1996
EP 0779291	A	18-06-1997	CA 2192281 A	12-06-1997
			JP 9202785 A	05-08-1997
			US 5922740 A	13-07-1999
WO 9802440	A	22-01-1998	AU 3441197 A	09-02-1998
			EP 0912578 A	06-05-1999
EP 0146243	A	26-06-1985	CA 1281329 A	12-03-1991
			JP 60112783 A	19-06-1985
			US 4745127 A	17-05-1988
			US 4822803 A	18-04-1989
			US 4933351 A	12-06-1990
			US 4663347 A	05-05-1987
WO 9802440	A	22-01-1998	AU 3441197 A	09-02-1998
			EP 0912578 A	06-05-1999
EP 0731099	A	11-09-1996	AT 193704 T	15-06-2000
			AU 705667 B	27-05-1999
			AU 4582896 A	19-09-1996
			BG 100396 A	31-12-1996
			BR 9600930 A	30-12-1997
			CA 2170824 A	07-09-1996
			CN 1134419 A	30-10-1996
			CZ 9600667 A	11-09-1996
			DE 69608721 D	13-07-2000
			FI 960998 A	07-09-1996
			HR 960071 A	31-12-1997
			HU 9600549 A	28-07-1997
			JP 8253466 A	01-10-1996
			NO 960884 A	09-09-1996
			NZ 286107 A	26-08-1998
			PL 313070 A	16-09-1996
			SK 29796 A	01-10-1996
			TR 960863 A	21-10-1996
			ZA 9601774 A	17-09-1996